

RUSSIAN ACADEMY OF SCIENCES
INSTITUTE FOR PROBLEMS IN MECHANICAL ENGINEERING
PETER THE GREAT ST. PETERSBURG POLYTECHNIC UNIVERSITY

Proceedings of the XLIV Summer School – Conference

ADVANCED PROBLEMS IN MECHANICS

A P M 2 0 1 6

St. Petersburg



POLYTECH

Peter the Great
St. Petersburg Polytechnic
University



IPME RAS

Institute for Problems in
Mechanical Engineering
Russian Academy of Sciences

<http://apm-conf.spb.ru>

St. Petersburg, 2016

SCIENTIFIC COMMITTEE AND EDITORIAL BOARD

- **D. A. Indeitsev** (*Co-Chairman*), IPME RAS, Peter the Great St. Petersburg Polytechnic University, Russia
- **A. M. Krivtsov** (*Co-Chairman*), Peter the Great St. Petersburg Polytechnic University, IPME RAS, Russia
- **P. A. Dyatlova** (*Scientific Secretary*), Peter the Great St. Petersburg Polytechnic University, IPME RAS, Russia
- **O. S. Loboda**, Peter the Great St. Petersburg Polytechnic University, IPME RAS, Russia
- **H. Altenbach**, Otto-von-Guericke University Magdeburg, Germany
- **V. A. Babeshko**, Southern Scientific Center RAS, Rostov-on-Don, Russia
- **A. K. Belyaev**, IPME RAS, St. Petersburg, Russia
- **I. E. Berinskii**, Tel Aviv University, Israel
- **I. I. Blekhman**, IPME RAS, Mekhanobr-tekhnika Corp., St. Petersburg, Russia
- **A. A. Burenin**, Institute of Metallurgy and Mechanical Engineering Far-Eastern Branch RAS, Komsomolsk-na-Amure, Russia
- **A. Castellanos**, University of Seville, Spain
- **A. V. Cherkaev**, University of Utah, Salt Lake City, USA
- **V. A. Eremeyev**, Rzeszow University of Technology, Poland
- **A. B. Freidin**, IPME RAS, Peter the Great St. Petersburg Polytechnic University, Russia
- **S. N. Gavrilov**, IPME RAS, St. Petersburg, Russia
- **R. V. Goldstein**, Institute for Problems in Mechanics RAS, Moscow, Russia
- **I. G. Goryacheva**, Institute for Problems in Mechanics RAS, Moscow, Russia
- **E. F. Grekova**, IPME RAS, St. Petersburg, Russia; University of Seville, Spain
- **N. Gupta**, Indian Institute of Technology, Delhi, India
- **H. Irschik**, Johannes Kepler University of Linz, Austria
- **M. L. Kachanov**, Tufts University, Medford, USA
- **B. L. Karihaloo**, Cardiff University, UK
- **V. A. Levin**, M. V. Lomonosov Moscow State University, Russia
- **A. M. Linkov**, IPME RAS, St. Petersburg, Russia; Rzeszow University of Technology, Poland
- **I. I. Lipatov**, Moscow Institute of Physics and Technology, Russia
- **E. V. Lomakin**, M. V. Lomonosov Moscow State University, Russia
- **A. V. Manzhurov**, Institute for Problems in Mechanics RAS, Moscow, Russia
- **G. A. Maugin**, Pierre and Marie Curie University, Paris, France
- **A. V. Metrikine**, Delft University of Technology, The Netherlands
- **G. Mishuris**, Aberystwyth University, UK
- **N. F. Morozov**, St. Petersburg State University, IPME RAS, Russia
- **W. H. Müller**, Technical University of Berlin, Germany
- **U. Nackenhorst**, Leibniz University of Hanover, Germany
- **V. A. Palmov**, Peter the Great St. Petersburg Polytechnic University, IPME RAS, Russia
- **E. Pavlovskaya**, University of Aberdeen, UK
- **Y. V. Petrov**, St. Petersburg State University, IPME RAS, Russia
- **M. B. Rubin**, Israel Institute of Technology, Haifa, Israel
- **A. I. Rudskoy**, Peter the Great St. Petersburg Polytechnic University, Russia
- **S. H. Sargsyan**, Gyumri State Pedagogical Institute, Armenia
- **E. N. Vilchevskaya**, Peter the Great St. Petersburg Polytechnic University, IPME RAS, Russia
- **M. Wiercigroch**, University of Aberdeen, UK
- **M. V. Zakrzhevsky**, Institute of Mechanics, Riga Technical University, Latvia

PREFACE

Dear Reader,

in this book you will find the Proceedings of the Summer School – Conference “Advanced Problems in Mechanics (APM) 2016”. The conference had been started in 1971. The first Summer School was organized by Prof. Ya.G. Panovko and his colleagues. In the early years the main focus of the School was on nonlinear oscillations of mechanical systems with a finite number of degrees of freedom. Since 1994 the Institute for Problems in Mechanical Engineering of the Russian Academy of Sciences organizes the Summer School. The traditional name of “Summer School” has been kept, but the topics covered by the School have been much widened, and the School has been transformed into an international conference. Now it is held under the patronage of the Russian Academy of Sciences. The topics of the conference cover now almost all fields of mechanics, being concentrated around the following main scientific directions:

- aerospace mechanics;
- computational mechanics;
- dynamics of rigid bodies and multibody dynamics;
- fluid and gas;
- mechanical and civil engineering applications;
- mechanics of media with microstructure;
- mechanics of granular media;
- nanomechanics;
- nonlinear dynamics, chaos and vibration;
- molecular and particle dynamics;
- phase transitions;
- solids and structures;
- wave motion.

The Summer School – Conference has two main purposes: to gather specialists from different branches of mechanics to provide a platform for cross-fertilization of ideas, and to give the young scientists a possibility to learn from their colleagues and to present their work. Thus the Scientific Committee encouraged the participation of young researchers, and did its best to gather at the conference leading scientists belonging to various scientific schools of the world.

We believe that the significance of Mechanics as of fundamental and applied science should much increase in the eyes of the world scientific community, and we hope that APM conference makes its contribution into this process.

We are happy to express our sincere gratitude for a partial financial support to Russian Foundation for Basic Research, Russian Academy of Sciences, and St. Petersburg Scientific Center. This support has helped substantially to organize the conference and to increase the participation of young researchers.

We hope that you will find the materials of the conference interesting, and we cordially invite you to participate in the coming APM conferences. You may find the information on the future “Advanced Problems in Mechanics” Schools – Conferences at our website:

<http://apm-conf.spb.ru>

With kind regards,

Co-Chairmen of APM 2016

Dmitri A. Indeitsev, Anton M. Krivtsov

Cases of integrability corresponding to the motion of a pendulum in the three-dimensional space

Maxim V. Shamolin
shamolin@rambler.ru, shamolin@imec.msu.ru

Abstract

In this activity, we systematize some results on the study of the equations of spatial motion of dynamically symmetric fixed rigid bodies–pendulums located in a nonconservative force fields. The form of these equations is taken from the dynamics of real fixed rigid bodies placed in a homogeneous flow of a medium. In parallel, we study the problem of a spatial motion of a free rigid body also located in a similar force fields. Herewith, this free rigid body is influenced by a nonconservative tracing force; under action of this force, either the magnitude of the velocity of some characteristic point of the body remains constant, which means that the system possesses a nonintegrable servo constraint, or the center of mass of the body moves rectilinearly and uniformly; this means that there exists a nonconservative couple of forces in the system.

1 Introduction

Earlier (see [1, 2]), the author already proved the complete integrability of the equations of a plane-parallel motion of a fixed rigid body–pendulum in a homogeneous flow of a medium under the jet flow conditions when the system of dynamical equations possesses a first integral, which is a transcendental (in the sense of the theory of functions of a complex variable, i.e., it has essential singularities) function of quasi-velocities. In [1, 2, 3], the planar problem was generalized to the spatial (three-dimensional) case, where the system of dynamical equations has a complete set of transcendental first integrals. It was assumed that the interaction of the homogeneous medium flow with the fixed body (the spherical pendulum) is concentrated on a part of the body surface that has the form of a planar (two-dimensional) disk. In this activity, the results relate to the case where all interaction of the homogeneous flow of a medium with the fixed body is concentrated on that part of the surface of the body, which has the form of a two-dimensional disk, and the action of the force is concentrated in a direction perpendicular to this disk. These results are systematized and are presented in invariant form.

2 Model assumptions

Let consider the homogeneous plane circle disk \mathcal{D} (with the center in the point D), the plane of which perpendicular to the holder OD . The disk is rigidly fixed perpendicular to the tool holder OD located on the spherical hinge O , and it flows about homogeneous fluid flow (Fig. 1). In this case, the body is a physical (spherical) pendulum. The medium flow moves from infinity with constant velocity $\mathbf{v} = \mathbf{v}_\infty \neq \mathbf{0}$. Assume that the holder does not create a resistance.




Figure 1: Fixed pendulum on a spherical hinge in the stream running medium

We suppose that the total force \mathbf{S} of medium flow interaction is parallel to the holder, and point N of application of this force is determined by at least the angle of attack α , which is made by the velocity vector \mathbf{v}_D of the point D with respect to the flow and the holder OD (Fig. 1); the total force is also determined by the angle β_1 , which is made in the plane of the disk \mathcal{D} (thus, (ν, α, β_1) are the spherical coordinates of the tip of the vector \mathbf{v}_D), and also the reduced angular velocity $\omega \cong l\Omega/v_D$, $v_D = |\mathbf{v}_D|$ (l is the length of the holder, Ω is the angular velocity of the pendulum). Such conditions arise when one uses the model of streamline flow around spatial bodies. Therefore, the force \mathbf{S} is directed along the normal to the disk to its side, which is opposite to the direction of the velocity \mathbf{v}_D , and passes through a certain point N of the disk such that the velocity vector \mathbf{v}_D and the force of the interaction \mathbf{S} lie in the plane ODN .

The vector $\mathbf{e} = \mathbf{OD}/l$ determines the orientation of the holder. Then $\mathbf{S} = s(\alpha)v_D^2\mathbf{e}$, $s(\alpha) = s_1(\alpha)\text{sign} \cos \alpha$, and the resistance coefficient $s_1 \geq 0$ depends only on the angle of attack α . By the axe-symmetry properties of the body–pendulum with respect to the point D , the function $s(\alpha)$ is even.

Let $Dx_1x_2x_3 = Dxyz$ be the coordinate system rigidly attached to the body, here-with, the axis $Dx = Dx_1$ has a direction vector \mathbf{e} , and the axes $Dx_2 = Dy$ and $Dx_3 = Dz$ lie in the plane of the disk \mathcal{D} (Fig. 1). In the same figure it is shown the angles $\theta = \xi$, $\psi = \eta_1$, i.e., the angles determining the pendulum position on the sphere. In this case, the angle θ is made by the holder and the direction of the over-running medium flow (the axis x_0); and the angle ψ is made by the projection

of the holder to the immovable plane y_0z_0 (which perpendicular to the over-running medium flow) and the axis y_0 (Fig. 1). Obviously, the angles $(\theta, \psi) = (\xi, \eta_1)$ are the spherical coordinates of the point D .

The space of positions of this spherical (physical) pendulum is the two-dimensional sphere

$$\mathbf{S}^2\{(\xi, \eta_1) \in \mathbf{R}^2 : 0 \leq \xi \leq \pi, \eta_1 \bmod 2\pi\}, \quad (1)$$

and its phase space is the tangent bundle of the two-dimensional sphere

$$T_*\mathbf{S}^2\{(\dot{\xi}, \dot{\eta}_1; \xi, \eta_1) \in \mathbf{R}^4 : 0 \leq \xi \leq \pi, \eta_1 \bmod 2\pi\}. \quad (2)$$

To the angular velocity, we put in correspondence $\Omega = \Omega_1\mathbf{e}_1 + \Omega_2\mathbf{e}_2 + \Omega_3\mathbf{e}_3$ ($\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$ the unit vectors of the coordinate system $Dx_1x_2x_3$) the skew-symmetric matrix

$$\tilde{\Omega} = \begin{pmatrix} 0 & -\Omega_3 & \Omega_2 \\ \Omega_3 & 0 & -\Omega_1 \\ -\Omega_2 & \Omega_1 & 0 \end{pmatrix}, \quad \tilde{\Omega} \in \mathfrak{so}(3). \quad (3)$$

The distance from the center D of the disk to the center of pressure (the point N , Fig. 1) has the form $|\mathbf{r}_N| = r_N = DN(\alpha, \beta_1, l\Omega/v_D)$, where $\mathbf{r}_N = \{0, x_{2N}, x_{3N}\} = \{0, y_N, z_N\}$ in system $Dx_1x_2x_3 = Dxyz$ (we omit the wave over Ω).

3 Set of dynamical equations in Lie algebra $\mathfrak{so}(3)$

If $\text{diag}\{I_1, I_2, I_2\}$ is the tensor of inertia of the body-pendulum in the coordinate system $Dx_1x_2x_3$ then the general equation of its motion has the following form:

$$\begin{aligned} I_1\dot{\Omega}_1 &= 0, \\ I_2\dot{\Omega}_2 + (I_1 - I_2)\Omega_1\Omega_3 &= -z_N \left(\alpha, \beta_1, \frac{\Omega}{v_D} \right) s(\alpha)v_D^2, \\ I_2\dot{\Omega}_3 + (I_2 - I_1)\Omega_1\Omega_2 &= y_N \left(\alpha, \beta_1, \frac{\Omega}{v_D} \right) s(\alpha)v_D^2, \end{aligned} \quad (4)$$

since the moment of the medium interaction force is determined by the following auxiliary matrix:

$$\begin{pmatrix} 0 & x_{2N} & x_{3N} \\ -s(\alpha)v_D^2 & 0 & 0 \end{pmatrix},$$

where $\{-s(\alpha)v_D^2, 0, 0\}$ is the decomposition of the medium interaction force \mathbf{S} in the coordinate system $Dx_1x_2x_3$.

Since the dimension of the Lie algebra $\mathfrak{so}(3)$ is equal to 3, the system of equations (4) is a group of dynamical equations on $\mathfrak{so}(3)$, and, simply speaking, the motion equations.

We see, that in the right-hand side of Eq. (4), first of all, it includes the angles α, β_1 , therefore, this system of equations is not closed. In order to obtain a complete system of equations of motion of the pendulum, it is necessary to attach several sets of kinematic equations to the dynamic equation on the Lie algebra $\mathfrak{so}(3)$.

3.1 Cyclic first integral

We immediately note that the system (4), by the existing dynamic symmetry

$$I_2 = I_3, \quad (5)$$

possesses the cyclic first integral

$$\Omega_1 \equiv \Omega_1^0 = \text{const.} \quad (6)$$

In this case, further, we consider the dynamics of our system at zero level:

$$\Omega_1^0 = 0. \quad (7)$$

Under conditions (5)–(7), the system (4) has the form of unclosed system of two equations:

$$I_2 \dot{\Omega}_2 = -z_N \left(\alpha, \beta_1, \frac{\Omega}{v_D} \right) s(\alpha) v_D^2, \quad I_2 \dot{\Omega}_3 = y_N \left(\alpha, \beta_1, \frac{\Omega}{v_D} \right) s(\alpha) v_D^2. \quad (8)$$

4 First set of kinematic equations

In order to obtain a complete system of equations of motion, it needs the set of kinematic equations which relate the velocities of the point D (i.e., the formal center of the disk D) and the over-running medium flow:

$$\mathbf{v}_D = v_D \cdot \mathbf{i}_v(\alpha, \beta_1) = \tilde{\Omega} \begin{pmatrix} l \\ 0 \\ 0 \end{pmatrix} + (-v_\infty) \mathbf{i}_v(-\xi, \eta_1), \quad (9)$$

$$\mathbf{i}_v(\alpha, \beta_1) = \begin{pmatrix} \cos \alpha \\ \sin \alpha \cos \beta_1 \\ \sin \alpha \sin \beta_1 \end{pmatrix}. \quad (10)$$

The equation (9) expresses the theorem of addition of velocities in projections on the related coordinate system $Dx_1x_2x_3$.

Indeed, the left-hand side of Eq. (9) is the velocity of the point D of the pendulum with respect to the flow in the projections on the related with the pendulum coordinate system $Dx_1x_2x_3$. Herewith, the vector $\mathbf{i}_v(\alpha, \beta_1)$ is the unit vector along the axis of the vector \mathbf{v}_D . The vector $\mathbf{i}_v(\alpha, \beta_1)$ has the spherical coordinates $(1, \alpha, \beta_1)$, which determines the decomposition (10).

The right-hand side of the Eq. (9) is the sum of the velocities of the point D when you rotate the pendulum (the first term), and the motion of the flow (the second term). In this case, in the first term, we have the coordinates of the vector $\mathbf{OD} = \{l, 0, 0\}$ in the coordinate system $Dx_1x_2x_3$.

We explain the second term of the right-hand side of Eq. (9) in more detail. We have in it the coordinates of the vector $(-\mathbf{v}_\infty) = \{-v_\infty, 0, 0\}$ in the immovable space. In order to describe it in the projections on the related coordinate system $Dx_1x_2x_3$, we need to make a (reverse) rotation of the pendulum at the angle $(-\xi)$ that is algebraically equivalent to multiplying the value $(-v_\infty)$ on the vector $\mathbf{i}_v(-\xi, \eta_1)$.

Thus, the first set of kinematic equations (9) has the following form in our case:

$$\begin{aligned} v_D \cos \alpha &= -v_\infty \cos \xi, \\ v_D \sin \alpha \cos \beta_1 &= l\Omega_3 + v_\infty \sin \xi \cos \eta_1, \\ v_D \sin \alpha \sin \beta_1 &= -l\Omega_2 + v_\infty \sin \xi \sin \eta_1. \end{aligned} \quad (11)$$

5 Second set of kinematic equations

We also need a set of kinematic equations which relate the angular velocity tensor $\tilde{\Omega}$ and coordinates $\xi, \eta_1, \dot{\xi}, \dot{\eta}_1$ of the phase space (2) of pendulum studied, i.e., the tangent bundle $T_*\mathbf{S}^2\{\xi, \eta_1; \dot{\xi}, \dot{\eta}_1\}$.

We draw the reasoning style allowing arbitrary dimension. The desired equations are obtained from the following two sets of relations. Since the motion of the body takes place in a Euclidean space $\mathbf{E}^n, n = 3$ formally, at the beginning, we express the tuple consisting of a phase variables Ω_2, Ω_3 , through new variable z_1, z_2 (from the tuple z). For this, we draw the following turn by the angle η_1 :

$$\begin{aligned} \begin{pmatrix} \Omega_2 \\ \Omega_3 \end{pmatrix} &= T_{1,2}(\eta_1) \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}, \\ T_{1,2}(\eta_1) &= \begin{pmatrix} \cos \eta_1 & -\sin \eta_1 \\ \sin \eta_1 & \cos \eta_1 \end{pmatrix}. \end{aligned} \quad (12)$$

In other words, the relations

$$\begin{pmatrix} z_1 \\ z_2 \end{pmatrix} = T_{1,2}(-\eta_1) \begin{pmatrix} \Omega_2 \\ \Omega_3 \end{pmatrix} \quad (13)$$

hold, i.e.,

$$z_1 = \Omega_2 \cos \eta_1 + \Omega_3 \sin \eta_1, \quad z_2 = -\Omega_2 \sin \eta_1 + \Omega_3 \cos \eta_1. \quad (14)$$

Then we substitute the following relationship instead of the variable z :

$$z_2 = \dot{\xi}, \quad z_1 = -\dot{\eta}_1 \frac{\sin \xi}{\cos \xi}. \quad (15)$$

Thus, two sets of Eqs. (12) and (15) give the second set of kinematic equations:

$$\Omega_2 = -\dot{\xi} \sin \eta_1 - \dot{\eta}_1 \frac{\sin \xi}{\cos \xi} \cos \eta_1, \quad \Omega_3 = \dot{\xi} \cos \eta_1 - \dot{\eta}_1 \frac{\sin \xi}{\cos \xi} \sin \eta_1. \quad (16)$$

We see that three sets of the relations (8), (11), and (16) form the closed system of equations.

These three sets of equations include the following functions:

$$y_N \left(\alpha, \beta_1, \frac{\Omega}{v_D} \right), \quad z_N \left(\alpha, \beta_1, \frac{\Omega}{v_D} \right), \quad s(\alpha). \quad (17)$$

In this case, the function s is considered to be dependent only on α , and the functions y_N, z_N may depend on, along with the angles α, β_1 , generally speaking, the reduced angular velocity $\omega \cong l\Omega/v_D$.

6 Problem on free body motion under assumption of tracing force

Parallel to the present problem of the motion of the fixed body, we study the spatial motion of the free axially symmetric rigid body with the frontal plane butt-end (the circle disk \mathcal{D}) in the resistance force fields under the quasi-stationarity conditions with the same model of medium interaction.

If (v, α, β_1) are the spherical coordinates of the velocity vector of the center D of disk \mathcal{D} lying on the axis of symmetry of a body, $\Omega = \{\Omega_1, \Omega_2, \Omega_3\}$ are the projections of its angular velocity on the axes of the coordinate system $Dx_1x_2x_3$ related to the body (in this case, the axis of symmetry CD coincides with the axis $Dx_1 = Dx$, C is the center of mass), and the axes $Dx_2 = Dy$ and $Dx_3 = Dz$ lie in the hyperplane of the disk; $I_1, I_2, I_3 = I_2, m$ are characteristics of inertia and mass, then the dynamical part of the equations of motion in which the tangent forces of the interaction of the body with the medium are absent, has the form

$$\begin{aligned} \dot{v} \cos \alpha - \dot{\alpha} v \sin \alpha + \Omega_2 v \sin \alpha \sin \beta_1 - \Omega_3 v \sin \alpha \cos \beta_1 + \sigma(\Omega_2^2 + \Omega_3^2) &= \frac{F_x}{m}, \\ \dot{v} \sin \alpha \cos \beta_1 + \dot{\alpha} v \cos \alpha \cos \beta_1 - \dot{\beta}_1 v \sin \alpha \sin \beta_1 + \Omega_3 v \cos \alpha - \\ - \Omega_1 v \sin \alpha \sin \beta_1 - \sigma \Omega_1 \Omega_2 - \sigma \dot{\Omega}_3 &= 0, \\ \dot{v} \sin \alpha \sin \beta_1 + \dot{\alpha} v \cos \alpha \sin \beta_1 + \dot{\beta}_1 v \sin \alpha \cos \beta_1 + \Omega_1 v \sin \alpha \cos \beta_1 - \\ - \Omega_2 v \cos \alpha - \sigma \Omega_1 \Omega_3 + \sigma \dot{\Omega}_2 &= 0, \\ I_1 \dot{\Omega}_1 &= 0, \\ I_2 \dot{\Omega}_2 + (I_1 - I_2) \Omega_1 \Omega_3 &= -z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) s(\alpha) v^2, \\ I_2 \dot{\Omega}_3 + (I_2 - I_1) \Omega_1 \Omega_2 &= y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) s(\alpha) v^2, \end{aligned} \quad (18)$$

$F_x = -S$, $S = s(\alpha)v^2$, $\sigma = CD$, in this case

$$\left(0, y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right), z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \right)$$

are the coordinates of the point N of application of the force \mathbf{S} in the coordinate system $Dx_1x_2x_3 = Dxyz$ related to the body.

The first part of three equations of the system (18) describe the motion of the center of a mass in the three-dimensional Euclidean space \mathbf{E}^3 in the projections on the coordinate system $Dx_1x_2x_3$. And the second part of three equation of the system (18) is obtained from the theorem on the change of the angular moment of a rigid body in the König axis.

Thus, the direct product $\mathbf{R}^1 \times \mathbf{S}^2 \times \mathfrak{so}(3)$ of the three-dimensional manifold and the Lie algebra $\mathfrak{so}(3)$ is the phase space of sixth-order system (18) of the dynamical equations. Herewith, since the medium influence force does not depend on the position of the body in a plane, the system (18) of the dynamical equations *is separated from the system of kinematic equations* and may be studied independently.

6.1 Cyclic first integral

We immediately note that the system (18), by the existing dynamic symmetry

$$I_2 = I_3, \quad (19)$$

possesses the cyclic first integral

$$\Omega_1 \equiv \Omega_1^0 = \text{const.} \quad (20)$$

In this case, further, we consider the dynamics of our system at zero level:

$$\Omega_1^0 = 0. \quad (21)$$

6.2 Nonintegrable constraint

If we consider *a more general problem* on the motion of a body under the action of a certain tracing force \mathbf{T} passing through the center of mass and providing the fulfillment of the equality

$$v \equiv \text{const}, \quad (22)$$

during the motion, then F_x in system (18) must be replaced by

$$T - s(\alpha)v^2. \quad (23)$$

As a result of an appropriate choice of the magnitude T of the tracing force, we can achieve the fulfillment of Eq. (22) during the motion. Indeed, if we formally express the value T by virtue of system (18), we obtain (for $\cos \alpha \neq 0$):

$$T = T_v(\alpha, \beta_1, \Omega) = m\sigma(\Omega_2^2 + \Omega_3^2) + \\ + s(\alpha)v^2 \left[1 - \frac{m\sigma \sin \alpha}{I_2 \cos \alpha} \left[z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \sin \beta_1 + y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \cos \beta_1 \right] \right]. \quad (24)$$

This procedure can be viewed from two standpoints. First, a transformation of the system has occurred at the presence of the tracing (control) force in the system which provides the corresponding class of motions (22). Second, we can consider this procedure as a procedure that allows one to reduce the order of the system. Indeed, system (18) generates an independent fourth-order system of the following form:

$$\begin{aligned} \dot{\alpha}v \cos \alpha \cos \beta_1 - \dot{\beta}_1 v \sin \alpha \sin \beta_1 + \Omega_3 v \cos \alpha - \sigma \dot{\Omega}_3 &= 0, \\ \dot{\alpha}v \cos \alpha \sin \beta_1 + \dot{\beta}_1 v \sin \alpha \cos \beta_1 - \Omega_2 v \cos \alpha + \sigma \dot{\Omega}_2 &= 0, \\ I_2 \dot{\Omega}_2 &= -z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) s(\alpha)v^2, \\ I_2 \dot{\Omega}_3 &= y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) s(\alpha)v^2, \end{aligned} \quad (25)$$

where the parameter v is supplemented by the constant parameters specified above.

The system (25) is equivalent to the system

$$\begin{aligned}
 \dot{\alpha}v \cos \alpha + v \cos \alpha [\Omega_3 \cos \beta_1 - \Omega_2 \sin \beta_1] + \sigma [-\dot{\Omega}_3 \cos \beta_1 + \dot{\Omega}_2 \sin \beta_1] &= 0, \\
 \dot{\beta}_1 v \sin \alpha - v \cos \alpha [\Omega_2 \cos \beta_1 + \Omega_3 \sin \beta_1] + \sigma [\dot{\Omega}_2 \cos \beta_1 + \dot{\Omega}_3 \sin \beta_1] &= 0, \\
 \dot{\Omega}_2 &= -\frac{v^2}{I_2} z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) s(\alpha), \\
 \dot{\Omega}_3 &= \frac{v^2}{I_2} y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) s(\alpha).
 \end{aligned} \tag{26}$$

We introduce new quasi-velocities in our system:

$$\begin{aligned}
 \begin{pmatrix} \Omega_2 \\ \Omega_3 \end{pmatrix} &= T_{1,2}(\beta_1) \begin{pmatrix} z_1 \\ z_2 \end{pmatrix}, \\
 T_{1,2}(\beta_1) &= \begin{pmatrix} \cos \beta_1 & -\sin \beta_1 \\ \sin \beta_1 & \cos \beta_1 \end{pmatrix}.
 \end{aligned} \tag{27}$$

In other words, the following relations

$$\begin{pmatrix} z_1 \\ z_2 \end{pmatrix} = T_{1,2}(-\beta_1) \begin{pmatrix} \Omega_2 \\ \Omega_3 \end{pmatrix} \tag{28}$$

hold, i.e.,

$$z_1 = \Omega_2 \cos \beta_1 + \Omega_3 \sin \beta_1, \quad z_2 = -\Omega_2 \sin \beta_1 + \Omega_3 \cos \beta_1. \tag{29}$$

We can see from (26) that the system cannot be solved uniquely with respect to $\dot{\alpha}$, $\dot{\beta}_1$ on the manifold

$$O = \left\{ (\alpha, \beta_1, \Omega_2, \Omega_3) \in \mathbf{R}^4 : \alpha = \frac{\pi}{2}k, k \in \mathbf{Z} \right\}. \tag{30}$$

Thus, formally speaking, the uniqueness theorem is violated on manifold (30). Moreover, the indefiniteness occurs for even k because of the degeneration of the spherical coordinates (v, α, β_1) , and an obvious violation of the uniqueness theorem for odd k occurs since the first equation of (26) is degenerate for this case.

This implies that system (25) outside of the manifold (30) (and only outside it) is equivalent to the following system:

$$\begin{aligned}
 \dot{\alpha} &= -z_2 + \frac{\sigma v s(\alpha)}{I_2 \cos \alpha} \left[z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \sin \beta_1 + y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \cos \beta_1 \right], \\
 \dot{z}_2 &= \frac{v^2}{I_2} s(\alpha) \left[z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \sin \beta_1 + y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \cos \beta_1 \right] - \\
 &- z_1^2 \frac{\cos \alpha}{\sin \alpha} - \frac{\sigma v s(\alpha)}{I_2 \sin \alpha} z_1 \left[z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \cos \beta_1 - y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \sin \beta_1 \right], \\
 \dot{z}_1 &= z_1 z_2 \frac{\cos \alpha}{\sin \alpha} + \left[-\frac{v^2}{I_2} s(\alpha) + \frac{\sigma v s(\alpha)}{I_2 \sin \alpha} z_2 \right] \times \\
 &\times \left[z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \cos \beta_1 - y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \sin \beta_1 \right], \\
 \dot{\beta}_1 &= z_1 \frac{\cos \alpha}{\sin \alpha} + \frac{\sigma v s(\alpha)}{I_2 \sin \alpha} \left[z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \cos \beta_1 - y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \sin \beta_1 \right].
 \end{aligned} \tag{31}$$

6.3 Constant velocity of the center of mass

If we consider *a more general problem* on the motion of a body under the action of a certain tracing force \mathbf{T} passing through the center of mass and providing the fulfillment of the equality

$$\mathbf{V}_C \equiv \text{const} \quad (32)$$

during the motion (\mathbf{V}_C is the velocity of the center of mass), then F_x in system (18) must be replaced by zero since the nonconservative couple of the forces acts on the body: $T - s(\alpha)v^2 \equiv 0$.

Obviously, we must choose the value of the tracing force T as follows:

$$T = T_v(\alpha, \beta_1, \Omega) = s(\alpha)v^2, \quad \mathbf{T} \equiv -\mathbf{S}. \quad (33)$$

The choice (33) of the magnitude of the tracing force T is a particular case of the possibility of separation of an independent fourth-order subsystem after a certain transformation of the system (18).

Indeed, let the following condition hold for T :

$$T = T_v(\alpha, \beta_1, \Omega) = \sum_{i,j=0, i \leq j}^3 \tau_{i,j} \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \Omega_i \Omega_j = T_1 \left(\alpha, \beta_1, \frac{\Omega}{v} \right) v^2, \quad \Omega_0 = v. \quad (34)$$

At the beginning, we introduce new quasi-velocities (27)–(29).

We rewrite the system (18) for the cases (19)–(21) in the form

$$\begin{aligned} & \dot{v} + \sigma(z_1^2 + z_2^2) \cos \alpha - \\ & - \sigma \frac{v^2}{I_2} s(\alpha) \sin \alpha \left[y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \cos \beta_1 + z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \sin \beta_1 \right] = \\ & = \frac{T_1 \left(\alpha, \beta_1, \frac{\Omega}{v} \right) v^2 - s(\alpha)v^2}{m} \cos \alpha, \\ & \dot{\alpha} v + z_2 v - \sigma(z_1^2 + z_2^2) \sin \alpha - \\ & - \sigma \frac{v^2}{I_2} s(\alpha) \cos \alpha \left[y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \cos \beta_1 + z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \sin \beta_1 \right] = \\ & = \frac{s(\alpha)v^2 - T_1 \left(\alpha, \beta_1, \frac{\Omega}{v} \right) v^2}{m} \sin \alpha, \\ & \dot{\Omega}_3 = \frac{v^2}{I_2} y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) s(\alpha), \\ & \dot{\Omega}_2 = -\frac{v^2}{I_2} z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) s(\alpha), \\ & \dot{\beta}_1 \sin \alpha - z_1 \cos \alpha - \\ & - \frac{\sigma v}{I_2} s(\alpha) \left[z_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \cos \beta_1 - y_N \left(\alpha, \beta_1, \frac{\Omega}{v} \right) \sin \beta_1 \right] = 0. \end{aligned} \quad (35)$$

If we introduce the new dimensionless phase variables and the differentiation by the formulas $z_k = n_1 v Z_k$, $k = 1, 2$, $\langle \cdot \rangle = n_1 v \langle' \rangle$, $n_1 > 0$, $n_1 = \text{const}$, system (35) has the following form:

$$v' = v \Psi(\alpha, \beta_1, Z_1, Z_2), \quad (36)$$

$$\begin{aligned} \alpha' = & -Z_2 + \sigma n_1 (Z_1^2 + Z_2^2) \sin \alpha + \\ & + \frac{\sigma}{I_2 n_1} s(\alpha) \cos \alpha [y_N(\alpha, \beta_1, n_1 Z) \cos \beta_1 + z_N(\alpha, \beta_1, n_1 Z) \sin \beta_1] - \\ & - \frac{T_1(\alpha, \beta_1, n_1 Z) - s(\alpha)}{m n_1} \sin \alpha, \end{aligned} \quad (37)$$

$$\begin{aligned} Z_2' = & \frac{s(\alpha)}{I_2 n_1^2} [1 - \sigma n_1 Z_2 \sin \alpha] [y_N(\alpha, \beta_1, n_1 Z) \cos \beta_1 + z_N(\alpha, \beta_1, n_1 Z) \sin \beta_1] - \\ & - \frac{\sigma}{I_2 n_1} Z_1 \frac{s(\alpha)}{\sin \alpha} [z_N(\alpha, \beta_1, n_1 Z) \cos \beta_1 - y_N(\alpha, \beta_1, n_1 Z) \sin \beta_1] - \\ & - Z_1^2 \frac{\cos \alpha}{\sin \alpha} + \sigma n_1 Z_2 (Z_1^2 + Z_2^2) \cos \alpha - Z_2 \frac{T_1(\alpha, \beta_1, n_1 Z) - s(\alpha)}{m n_1} \cos \alpha, \end{aligned} \quad (38)$$

$$\begin{aligned} Z_1' = & \frac{1}{I_2 n_1^2} \frac{s(\alpha)}{\sin \alpha} [\sigma n_1 Z_2 \sin \alpha - 1] [z_N(\alpha, \beta_1, n_1 Z) \cos \beta_1 - y_N(\alpha, \beta_1, n_1 Z) \sin \beta_1] - \\ & - \frac{\sigma}{I_2 n_1} Z_1 s(\alpha) \sin \alpha [z_N(\alpha, \beta_1, n_1 Z) \sin \beta_1 + y_N(\alpha, \beta_1, n_1 Z) \cos \beta_1] + \\ & + Z_1 Z_2 \frac{\cos \alpha}{\sin \alpha} + \sigma n_1 Z_1 (Z_1^2 + Z_2^2) \cos \alpha - Z_1 \frac{T_1(\alpha, \beta_1, n_1 Z) - s(\alpha)}{m n_1} \cos \alpha, \end{aligned} \quad (39)$$

$$\beta_1' = Z_1 \frac{\cos \alpha}{\sin \alpha} + \frac{\sigma}{I_2 n_1} \frac{s(\alpha)}{\sin \alpha} [z_N(\alpha, \beta_1, n_1 Z) \cos \beta_1 - y_N(\alpha, \beta_1, n_1 Z) \sin \beta_1], \quad (40)$$

$$\begin{aligned} \Psi(\alpha, \beta_1, Z_1, Z_2) = & -\sigma n_1 (Z_1^2 + Z_2^2) \cos \alpha + \\ & + \frac{\sigma}{I_2 n_1} s(\alpha) \sin \alpha [y_N(\alpha, \beta_1, n_1 Z) \cos \beta_1 + z_N(\alpha, \beta_1, n_1 Z) \sin \beta_1] + \\ & + \frac{T_1(\alpha, \beta_1, n_1 Z) - s(\alpha)}{m n_1} \cos \alpha. \end{aligned}$$

We see that the independent fourth-order subsystem (37)–(40) can be substituted into the fifth-order system (36)–(40) and can be considered separately on its own four-dimensional phase space.

7 Case where the moment of nonconservative forces is independent of the angular velocity

We take the function \mathbf{r}_N as follows (the disk \mathcal{D} is given by the equation $x_{1N} \equiv 0$):

$$\mathbf{r}_N = R(\alpha) \mathbf{i}_N, \quad (41)$$

where $\mathbf{i}_N = \mathbf{i}_v(\pi/2, \beta_1)$ (see (10)).

Thus, the equalities $x_{2N} = R(\alpha) \cos \beta_1$, $x_{3N} = R(\alpha) \sin \beta_1$ hold and show that for the considered system, the moment of the nonconservative forces is independent of the angular velocity (it depends only on the angles α, β_1).

And so, for the construction of the force field, we use the pair of dynamical functions $R(\alpha), s(\alpha)$; the information about them is of a qualitative nature. Similarly to the choice of the Chaplygin analytical functions, we take the dynamical functions s and R as follows:

$$R(\alpha) = A \sin \alpha, \quad s(\alpha) = B \cos \alpha, \quad A, B > 0. \quad (42)$$

Theorem 7.1. *The simultaneous equations (4), (11), (16), under conditions (5)–(7), (41), (42) can be reduced to the dynamical system on the tangent bundle (2) of the two-dimensional sphere (1).*

Indeed, if we introduce the dimensionless parameter and the differentiation by the formulas $b_* = ln_0, n_0^2 = AB/I_2, \langle \cdot \rangle = n_0 v_\infty \langle ' \rangle$, then the obtained equations have the following form:

$$\begin{aligned} \xi'' + b_* \xi' \cos \xi + \sin \xi \cos \xi - \eta_1'^2 \frac{\sin \xi}{\cos \xi} &= 0, \\ \eta_1'' + b_* \eta_1' \cos \xi + \xi' \eta_1' \frac{1 + \cos^2 \xi}{\cos \xi \sin \xi} &= 0, \quad b_* > 0. \end{aligned} \quad (43)$$

The phase pattern of the system (43) ($\xi \leftrightarrow \theta, \eta_1 \leftrightarrow \psi$) is shown in Fig. 2.




Figure 2: Phase pattern of the fixed pendulum on a spherical hinge situated in the medium flow

Theorem 7.2. *Three sets of relations (4), (11), (16) under conditions (5)–(7), (41), (42) possess three the first integrals (the complete set), which are the transcendental function (in the sense of complex analysis) and are expressed as a finite combination of elementary functions.*

We have the following topological and mechanical analogies in the sense explained above. (1) A motion of a fixed physical pendulum on a spherical hinge in a flowing medium (nonconservative force fields). (2) A spatial free motion of a rigid body in a nonconservative force field under a tracing force (in the presence of a nonintegrable constraint). (3) A spatial composite motion of a rigid body rotating about its center of mass, which moves rectilinearly and uniformly, in a nonconservative force field. On more general topological analogues, see also [4, 5, 6, 7, 8, 9].

Acknowledgements

This work was supported by the Russian Foundation for Basic Research, project no. 15-01-00848-a.

References

- [1] M. V. Shamolin, “Variety of Integrable Cases in Dynamics of Low- and Multi-Dimensional Rigid Bodies in Nonconservative Force Fields”, *Journal of Mathematical Sciences*, 204:4 (2015), 379–530.
- [2] M. V. Shamolin, “Dynamical Pendulum-Like Nonconservative Systems”, *Applied Non-Linear Dynamical Systems*, Springer Proceedings in Mathematics and Statistics, Vol. 93 (2014), 503–525.
- [3] M. V. Shamolin, “New Case of Integrability in the Dynamics of a Multidimensional Solid in a Nonconservative Field”, *Doklady Physics*, 58:11 (2013), 496–499.
- [4] M. V. Shamolin, “New case of integrability of dynamic equations on the tangent bundle of a 3-sphere”, *Russian Math. Surveys*, 68:5 (2013), 963–965.
- [5] M. V. Shamolin, “A New Case of Integrability in the Dynamics of a Multidimensional Solid in a Nonconservative Field under the Assumption of Linear Damping”, *Doklady Physics*, 59:8 (2014), 375–378.
- [6] M. V. Shamolin, “A Multidimensional Pendulum in a Nonconservative Force Field”, *Doklady Physics*, 60:1 (2015), 34–38.
- [7] M. V. Shamolin, “Classification of Integrable Cases in the Dynamics of a Four-Dimensional Rigid Body in a Nonconservative Field in the Presence of a Tracking Force”, *Journal of Mathematical Sciences*, 204:6 (2015), 808–870.
- [8] V. V. Trofimov, “Euler equations on Borel subalgebras of semisimple Lie algebras,” *Izv. Akad. Nauk SSSR, Ser. Mat.*, **43**, No. 3 (1979), 714–732.

REFERENCES

- [9] V. V. Trofimov, “Finite-dimensional representations of Lie algebras and completely integrable systems,” *Mat. Sb.*, **111**, No. 4 (1980), 610–621.

Maxim V. Shamolin, Lomonosov Moscow State University, Russian Federation