

Moscow Institute of Physics and Technology (Dolgoprudny)  
Steklov Mathematical Institute RAS (Moscow)  
Russian Foundation of Basic Research  
Journal "Regular and Chaotic Dynamics"



The International Scientific Workshop  
**"Recent Advances in Hamiltonian  
and Nonholonomic Dynamics"**

**Book of Abstracts**

15–18 June 2017  
Dolgoprudny, Russia

Moscow Institute of Physics and Tehnology (Dolgoprudny)  
Steklov Mathematical Institute RAS (Moscow)  
Russian Foundation of Basic Research  
Journal "Regular and Chaotic Dynamics"



The International Scientific Workshop  
**“Recent Advances in Hamiltonian  
and Nonholonomic Dynamics”**

**Book of Abstracts**

15–18 June 2017

Dolgoprudny, Russia

The International Scientific Workshop “Recent Advances in Hamiltonian and Nonholonomic Dynamics” is being held under the auspices of the Russian Foundation for Basic Research (17-01-20218).

**The International Scientific Workshop “Recent Advances in Hamiltonian and Nonholonomic Dynamics”** : Book of Abstracts. — Moscow–Izhevsk : Publishing Center “Institute of Computer Science”, 2017. — 88 p.

**ISBN 978-5-4344-0445-7**

## SCIENTIFIC COMMITTEE

### Chairs:

*Valery Kozlov*, V. A. Steklov Mathematical Institute of Russian Academy of Sciences, Russia

*Alexey V. Borisov*, Institute of Computer, Udmurt State University, Izhevsk, Russia

*Anastasios Bountis*, Nazarbayev University, Kazakhstan

*Alexander Kilin*, Udmurt State University, Izhevsk, Russia

*Alexander Ivanov*, Moscow Institute of Physics and Technology, Dolgoprudny, Russia

*Sergey Kuznetsov*, Kotel'nikov's Institute of Radio-Engineering and Electronics of RAS, Russia

*Ivan Mamaev*, Udmurt State University, Izhevsk, Russia

*Dmitry Treschev*, V. A. Steklov Mathematical Institute of Russian Academy of Sciences, Russia

## ORGANIZING COMMITTEE

Alexey Borisov, borisov@rcd.ru

Anastasios Bountis, tassosbountis@gmail.com

Alexander Kilin, aka@rcd.ru

Ivan Mamaev, mamaev@rcd.ru

Sergey Semendyaev, semendyaevsergey@gmail.com

Sergey Sokolov, sokolovsv72@mail.ru

# **Integrable systems with dissipation on the tangent bundle of a two-dimensional manifold**

Maxim V. Shamolin

*Lomonosov Moscow State University, Moscow, Russian Federation*

We study nonconservative systems for which the usual methods of study, e.g., Hamiltonian systems, are inapplicable. Thus, for such systems, we must “directly” integrate the main equation of dynamics. We generalize previously known cases and obtain new completely integrable cases in transcendental functions of the equation of dynamics of a four-dimensional rigid body in a nonconservative force field.

We obtain a series of complete integrable nonconservative dynamical systems with nontrivial symmetries. Moreover, in almost all cases, all first integrals are expressed in terms of finite combinations of elementary functions; these first integrals are transcendental functions of their variables. In this case, the transcendence is understood in the sense of complex analysis, when the analytic continuation of a function into the complex plane has essentially singular points. This fact is caused by the existence of attracting and repelling limit sets in the system (for example, attracting and repelling focuses).

We detect new integrable cases of the motion of a rigid body, including the classical problem of the motion of a multi-dimensional spherical pendulum in a flowing medium.

This activity is devoted to general aspects of the integrability of dynamical systems with variable dissipation. First, we propose a descriptive characteristic of such systems. The term “variable dissipation” refers to the possibility of alternation of its sign rather than to the value of the dissipation coefficient (therefore, it is more reasonable to use the term “sign-alternating”) [1, 2].

We introduce a class of autonomous dynamical systems with one periodic phase coordinate possessing certain symmetries that are typical of pendulum-type systems. We show that this class of systems can be naturally embedded in the class of systems with variable dissipation with zero mean, i.e., on the average for the period with respect to the periodic coordinate, the dissipation in the system is equal to zero, although in various domains of the phase space, either energy pumping or dissipation can occur, but they balance to each other in a certain sense. We present some examples of pendulum-type systems on lower-dimension manifolds from the dynamics of a rigid body in a nonconservative field.

Then we study certain general conditions of the integrability in elementary functions for systems on the two-dimensional plane and the tangent bundles

of a one-dimensional sphere (i.e., the two-dimensional cylinder) and a two-dimensional sphere (a four-dimensional manifold). Therefore, we propose an interesting example of a three-dimensional phase portrait of a pendulum-like system which describes the motion of a spherical pendulum in a flowing medium (see also [2, 3]).

To understand the difficulty of the problem to be solved, let us consider, for instance, the spherical pendulum ( $\psi$  and  $\theta$  are the coordinates of the point on the sphere where the pendulum is defined) in a jet flow. Then the equations of its motion are

$$\ddot{\theta} + (b_* - H_1^*)\dot{\theta} \cos \theta + \sin \theta \cos \theta - \dot{\psi}^2 \frac{\sin \theta}{\cos \theta} = 0, \quad (1)$$

$$\ddot{\psi} + (b_* - H_1^*)\dot{\psi} \cos \theta + \dot{\theta} \dot{\psi} \frac{1 + \cos^2 \theta}{\cos \theta \sin \theta} = 0, \quad b_* > 0, \quad H_1^* > 0, \quad (2)$$

and the phase pattern of eqs. (1), (2) is shown in Fig. 1.




Fig. 1. Phase pattern of spherical pendulum in a jet flow

The assertions obtained in the work for variable dissipation systems are a continuation of the Poincaré–Bendixon theory for systems on closed two-dimensional manifolds and the topological classification of such systems.

The problems considered in the work stimulate the development of qualitative tools of studying, and, therefore, in a natural way, there arises a qualitative variable dissipation system theory.

Following Poincaré, we improve some qualitative methods for finding key trajectories, i.e., the trajectories such that the global qualitative location of all

other trajectories depends on the location and the topological type of these trajectories. Therefore, we can naturally pass to a complete qualitative study of the dynamical system considered in the whole phase space. We also obtain a condition for the existence of bifurcation of the birth of stable and unstable limit cycles for the systems describing the body motion in a resisting medium under the streamline flow around. We find methods for finding any closed trajectories in the phase spaces of such systems and also present criteria for the absence of any such trajectories. We extend the Poincaré topographical plane system theory and the comparison system theory to the spatial case. We study some elements of the theory of monotone vector fields on orientable surfaces.

### References

- [1] Shamolin M. V. Dynamical Systems with Variable Dissipation: Approaches, Methods, and Applications// *Journal of Mathematical Sciences*, 2009, vol. 162, no. 6, pp. 741–908.
- [2] Trofimov V. V., Shamolin M. V. Geometric and Dynamical Invariants of Integrable Hamiltonian and Dissipative Systems// *Journal of Mathematical Sciences*, 2012, vol. 180, no. 4, pp. 365–530.
- [3] Shamolin M. V. Variety of Integrable Cases in Dynamics of Low- and Multi-Dimensional Rigid Bodies in Nonconservative Force Fields// *Journal of Mathematical Sciences*, 2015, vol. 203, no. 1, pp. 379–530.